


6



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search

PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Biochem J. 1996 Mar 1;314 (Pt 2):555-61.

Related Articles, Links

Biochem J

P2-purigenic receptors regulate phospholipase C and adenylate cyclase activities in immortalized Schwann cells.**Berti-Mattera LN, Wilkins PL, Madhun Z, Suchovsky D.**

Division of Hypertension, Department of Medicine, Case Western Reserve University School of Medicine and University Hospitals of Cleveland, OH 44106, USA.

Schwann cells play an important role in both the development and regeneration of peripheral nerves. Proliferation and differentiation of Schwann cells are critically dependent on changes in the levels of cAMP. ATP is a fast excitatory transmitter in the peripheral nervous system, inducing depolarization of the vagus nerve through occupancy of P2-purigenic receptors. In the present study we demonstrate that extracellular ATP stimulates phospholipase C and inhibits adenylate cyclase activities in cultured Schwann cells. Addition of ATP inhibited, in a concentration-dependent manner, forskolin- or isoprenaline-stimulated adenylate cyclase activity. The rank order of potency corresponding to different purinergic receptor agonists was 2-methylthio-ATP > ATP = ADP > or = adenosine 5'-[gamma-thio]triphosphate (ATP[S]) > UTP, consistent with the involvement of a P2y subtype. Adenosine and adenosine 5'-[alpha,beta-methylene]-triphosphate (pp[CH2pA]) were ineffective. Preincubation with pertussis toxin completely blocked this inhibitory effect. When Schwann cells were pre-labelled with myo-[3H]inositol and incubated in Hanks' balanced salt solution containing Ca²⁺ and Mg²⁺, addition of ATP[S] resulted in a concentration-dependent increase in the release of InsP with a concomitant increase in intracellular free [Ca²⁺] ([Ca²⁺]_i). Under these conditions, the effects of both ATP and UTP were of lower magnitude. Removal of Ca²⁺ and Mg²⁺ from the assay medium resulted in a significant increase in the effects of ATP[S], ATP and UTP. The decreased response observed in the presence of both bivalent cations (1.2 mM Ca²⁺ and 1 mM Mg²⁺) could not be explained either by increased degradation of ATP by Ca²⁺/Mg²⁺-dependent nucleotidases or by cation influx. The rank order of potency for the effects of agonists on phospholipase C activity was ATP[S] = adenosine 5'-[gamma-imido]triphosphate > ATP - UTP > ADP, indicating the involvement of a P(2U) receptor subtype in this response. Adenosine, AMP and pp[CH2]pA were ineffective. These results demonstrate that

immortalized Schwann cells express P(2U) and P(2Y) purinoceptors, which are coupled to stimulation of phospholipase C and inhibition of adenylate cyclase, respectively. Our observations unveil signal-transduction pathways that may be used by ATP to regulate proliferation and differentiation of Schwann cells, and ultimately to influence nerve homeostasis.

PMID: 8670070 [PubMed - indexed for MEDLINE]

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Jun 5 2003 10:08:34